

Overzicht meetinstrumenten NESDO

 = Key artikel niet aanwezig

Topic	Measurement instrument	Type*	Relevant references to literature
Screening			
- Screening for depression	Geriatric Depression Scale (GDS)	SR	Yesavage et al. 1982;17(1):37-49. Yesavage 1988; 24(4):709-11; Wancata et al. 2006;114(6):398-410.
- Telephone screening	Basic CIDI Questions for depression	INT	See CIDI
Demographics: Age, gender, education, SES, religion, partner status, family	Standard questions	INT	n/a**
Diagnostics	CIDI (life time): depression CIDI (12mnd): anxiety disorders	INT	Composite Interview Diagnostic Instrument (CIDI) –version 2.1 (World Health Organization, 1997) Andrews 1998;33(2):80-8; Wittchen et al 1989;46: 437-443; Wittchen et al. 1991;159:645-53, 658; Wittchen 1994;28(1):57-84.
Psychopathology			
- Depression, severity	Inventory of Depressive Symptoms (IDS)	SR2	Trivedi et al. 2004, 34, 73-82; Rush et al. 1996, 26:477-86
- Depression, symptoms	CES-d	SR1	Beekman et al. 1997;27(1):231-235; Radloff 1977; 3: 385-401
- Apathy, severity	Apathy Scale	SR1	 Starkstein 1992; 4(2):134-9
- Anxiety symptoms	Beck Anxiety Index	SR1	Beck et al. 1988;56(6):893-7. Steer et al. 1993;60(3):532-42. Kabacoff et al. 1997
Physical conditions			
- Chronic diseases	CBS/LASA Questionnaire	INT	Kriegsman 1996; 49:1407-1417
- Pain	Chronic Graded Pain Scale	INT	Von Korff 2005;117(3):304-13.
- Vision and hearing	LASA Questionnaires	INT	Pronk 2009 (documentatie LASA), van Nispen et al 2010 (in press)
- Sleep	Insomnia Rating Scale (IRS)	SR2	Levine et al. 2003 15: 137-148. Levine et al. 2003;15(2):123-36. Levine et al. 2005; 67(1):98-104
- Functional limitations	WHO-DAS II	SR2	Chwastiak et al. 2003,56: 507-514

- Medication use	Registration of medication use	INT	n/a
- Smoking & drugs	Past + current smoking questions	INT	n/a
- Alcohol dependency	AUDIT	SR2	Babor et al. 1989;14(2):139-57; Babor et al 2001 (guidelines)
- Physical Activities	International Physical Activities CBS	SR1	Craig et al. 2003 35: 1381-1395
- Mortality / cause of death		tzt	
Biological determinants			
- Metabolic syndrome, inflammation markers & polymorphism	Fasting blood sample	MED	n/a
- Autonomic Nervous System	Electro + impedance cardiography	MED	de Geus et al. 1995 41: 205-227
- Cortisol	6 saliva samples, including DST test	MED	n/a
Psychol. determinants			
- Personality: Big Five	NEO-FFI	SR1	Costa e & McCrae 1995 64: 21-50; Hoekstra et al. 1995 (Handleiding)
- Mastery	Pearlin Mastery scale	SR1	Pearlin & Schooler 1978, 19: 2-21.
- Daily Hassles	Worry Scale-R (15)	SR1	Raabe et al (interne rapportage); Wisocki PA 1986
Social determinants			
- Early life events	NEMESIS questionnaire	INT	De Graaf et al 2004 82(3):461-7.
- Recent life events	Brugha Questionnaire	INT	Brugha et al 1985 15: 189-194.
- Loneliness and affiliation	Loneliness and affiliation scale	SR1	De Jong Gierveld & Kamphuis 1995 9: 289-299; van Tilburg 1999 30(4):158-63
- Social support	Close Person Inventory (CPI)	SR2	Stansfeld et al. 1992 35: 1027-1035
Cognitive functioning			
- Global cogn. functioning	MMSE	INT	Folstein 1975 12(3), 189-198
- Executive functioning	STROOP (short form)	INT	Stroop 1935, 18:643-662 (n/a). Klein et al. 1997, 18(1): 77-82.
- Motivational problems	Digit Span (subtest WAIS)	INT	Wechsler , 1958 (n/a).
- Memory function	10-words test (adapted from Auditory Verbal Learning Test)	INT	Rey 1964 (n/a); van der Elst 2005 11(3):290-302
Use of Care	TIC-P	INT	Hakkaart-van Roijen, 2002

Evaluation - proxy information - permission further research		INT	
---	--	-----	--

* type of assessment: INT=interview, SR1=self-report questionnaire 1, SR2=self-report questionnaire 2, MED=Medical assessment

** n/a not available

Schriftelijke tussenmetingen (ieder half jaar):

- Veranderingen in wonen en gezondheid
- IDS (depressie ernst symptomen)
- WHO-DAS (Functionele beperkingen)
- BAI (angstsymptomen)
- TIC-P (Zorggebruik)
- Brugha (recente life events)
- Medicijngebruik

Reference List

- Andrews G, Peters L. The psychometric properties of the Composite International Diagnostic Interview. *Soc Psychiatry Psychiatr Epidemiol.* 1998;33(2):80-8. Review.
- Babor TF, Kranzler HR, Lauerman RJ. Early detection of harmful alcohol consumption: comparison of clinical, laboratory, and self-report screening procedures. *Addict Behav.* 1989;14(2):139-57.
- Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG. AUDIT. The alcohol use disorders identification test. Guidelines for use in Primary care. Second edition. World health organisation, 2001.
- Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol.* 1988;56(6):893-7.
- Beekman AT, Deeg DJ, Van Limbeek J, et al: Criterion validity of the Center for Epidemiologic Studies Depression scale (CES-D): results from a community-based sample of older subjects in The Netherlands. *Psychol Med* 1997;27(1):231-235
- Brugha T, Bebbington P, Tennant C, Hurry J. The List of Threatening Experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychol Med.* 1985, 15: 189-194.
- Chwastiak LA, Von Korff M. Disability in depression and back pain: evaluation of the World Health Organization Disability Assessment Schedule (WHO DAS II) in a primary care setting. *J Clin Epidemiol.* 2003, 56: 507-514.
- Costa PT, McCrae RR. Domains and facets: hierarchical personality assessment using the revised NEO personality inventory. *J Pers Assess.* 1995, 64: 21-50.
- Craig CL, Marshall AL, Sjostrom M, Bauman AE, Booth ML, Ainsworth BE, Pratt M, Ekelund U, Yngve A, Sallis JF, Oja P. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 35: 1381-1395.
- de Geus EJ, Willemsen GH, Klaver CH, van Doornen LJ (1995) Ambulatory measurement of respiratory sinus arrhythmia and respiration rate. *Biol Psychol.* 2003, 41: 205-227.
- de Graaf R, Bijl RV, Ten Have M, Beekman AT, Vollebergh WA. Pathways to comorbidity: the transition of pure mood, anxiety and substance use disorders into comorbid conditions in a longitudinal population-based study. *J Affect Disord.* 2004 82(3):461-7.
- DeJong Gierveld J, Kamphuis. The development of a rasch type loneliness scale. *Applied Psychological Measurement.* 1985, 9: 289-299.
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research,* 1975, 12(3), 189-198.
- Hakkaart-van Roijen L Trimbos/iMTA questionnaire for costs associated with psychiatric illness (TIC-P). Rotterdam: Institute for Medical Technology Assessment, 2002.

- Hoekstra HA, Ormel J, de Fruyt F. NEO-PR / NEO-FFI. Big Five Persoonlijkheidsvragenlijsten. Handleiding. Swets Test Services, Lisse, 1996.
- Klein M, Ponds RW, Houx PJ, Jolles J. Effect of test duration on age-related differences in Stroop interference. *J Clin Exp Neuropsychol.* 1997;19(1):77-82.
- Kriegsman DMW, Penninx BWJH, van Eijk JTM, et al: Self-reports and general practitioner information on the presence of chronic diseases in community dwelling elderly. A study on the accuracy of patients' self-reports and on determinants of inaccuracy. *J Clin Epidemiol* 1996; 49:1407-1417
- Levine DW, Kaplan RM, Kripke DF, Bowen DJ, Naughton MJ, Shumaker SA. Factor structure and measurement invariance of the Women's Health Initiative Insomnia Rating Scale. *Psychol Assess.* 2003;15(2):123-36.
- Levine DW, Kripke DF, Kaplan RM, Lewis MA, Naughton MJ, Bowen DJ, Shumaker SA. Reliability and validity of the Women's Health Initiative Insomnia Rating Scale. *Psychol Assess.* 2003;15(2):137-48.
- Levine DW, Dailey ME, Rockhill B, Tipping D, Naughton MJ, Shumaker SA. Validation of the Women's Health Initiative Insomnia Rating Scale in a multicenter controlled clinical trial. *Psychosom Med.* 2005;67(1):98-104.
- Pearlin LI, Schooler C. The structure of coping. *J Health Soc Behav.* 1978, 19: 2-21.
- Raabe S, Weijers A & Emmelkamp PMG. De ontwikkeling van een korte angstvragenlijst voor ouderen op basis van de Worry Scale en de Fear Questionnaire: eerste resultaten bij een niet-klinische Nederlandse populatie. Interne publicatie, UvA.
- Radloff LS: The CES-D Scale: A self-report depression scale for research in the general population. *Appl Psychol Meas.*1977; 3: 385-401
- Rey, A. L'examen clinique en psychologie (Clinical examination in psychology) Presse Universitaire de France, Paris, 1964.
- Rush AJ, Gullion CM, Basco MR, Jarrett RB, Trivedi MH. The Inventory of Depressive Symptomatology (IDS): psychometric properties. *Psychol Med.* 1996, 26: 477-486.
- Stansfeld S, Marmot M. Deriving a survey measure of social support: the reliability and validity of the Close Persons Questionnaire. *Soc Sci Med*, 1992, 35: 1027-1035.
- Starkstein SE, Mayberg HS, Preziosi TJ, Andrezejewski P, Leiguarda R, Robinson RG. Reliability, validity, and clinical correlates of apathy in Parkinson's disease. *J Neuropsychiatry Clin Neurosci.* 1992; 4(2):134-9.
- Steer RA, Rissmiller DJ, Ranieri WF, Beck AT. Structure of the computer-assisted Beck Anxiety Inventory with psychiatric inpatients. *J Pers Assess.* 1993, 60(3):532-42.
- Stroop JR. Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, 1935, 18:643-662

Trivedi MH, Rush AJ, Ibrahim HM, Carmody TJ, Biggs MM, Suppes T, Crismon ML, Shores-Wilson K, Toprac MG, Dennehy EB, Witte B, Kashner TM. The Inventory of Depressive Symptomatology, Clinician Rating (IDS-C) and Self-Report (IDS-SR), and the Quick Inventory of Depressive Symptomatology, Clinician Rating (QIDS-C) and Self-Report (QIDS-SR) in public sector patients with mood disorders: a psychometric evaluation. *Psychol Med.* 2004;34(1):73-82.

Van der Elst W, van Boxtel MP, van Breukelen GJ, Jolles J. Rey's verbal learning test: normative data for 1855 healthy participants aged 24-81 years and the influence of age, sex, education, and mode of presentation. *J Int Neuropsychol Soc.* 2005;11(3):290-302.

van Nispen RM, Knol DL, Mokkink LB, Comijs HC, Deeg DJ, van Rens GH. Vision-related quality of life Core Measure (VCM1) showed low-impact differential item functioning between groups with different administration modes. *J Clin Epidemiol.* 2010 in press

van Tilburg TG, de Jong Gierveld J. [Reference standards for the loneliness scale]. *Tijdschr Gerontol Geriatr.* 1999;30(4):158-63. Dutch.

Von Korff M, Miglioretti DL. A prognostic approach to defining chronic pain. *Pain.* 2005;117(3):304-13.

Wancata J, Alexandrowicz R, Marquart B, Weiss M, Friedrich F. The criterion validity of the Geriatric Depression Scale: a systematic review. *Acta Psychiatr Scand.* 2006;114(6):398-410.

Wechsler D. The measurement and appraisal of adult intelligence (4th. ed) Baltimore: Williams & Wilkins.

Wittchen HU. Reliability and validity studies of the WHO—Composite International Diagnostic Interview (CIDI): a critical review. *J Psychiatr Res.* 1994;28(1):57-84.

Wittchen HU, Burke JD, Semler G, Pfister H, Von CM, Zaudig M. Recall and dating of psychiatric symptoms. Test-retest reliability of time-related symptom questions in a standardized psychiatric interview. *Arch Gen Psychiatry.* 1989, 46: 437-443.

Wittchen HU, Robins LN, Cottler LB, Sartorius N, Burke JD, Regier D. Cross-cultural feasibility, reliability and sources of variance of the Composite International Diagnostic Interview (CIDI). The Multicentre WHO/ADAMHA Field Trials. *Br J Psychiatry,* 1991, 159: 645-53, 658.

Yesavage JA. Geriatric Depression Scale. *Psychopharmacol Bull.* 1988;24(4):709-11.

Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, Leirer VO. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res.* 1982-1983;17(1):37-49.